

VOTING IN AGREEABLE SOCIETIES

DEBORAH E. BERG, SERGUEI NORINE, FRANCIS EDWARD SU*, ROBIN THOMAS,
AND PAUL WOLLAN

My idea of an agreeable person... is a person who agrees with me.
—Benjamin Disraeli [8]

1. INTRODUCTION

When is agreement possible? An important aspect of group decision-making is the question of how a group makes a choice when individual preferences may differ. Clearly, when making a single group choice, people cannot all have their “ideal” preferences, i.e, the options that they most desire, if those ideal preferences are different. However, for the sake of agreement, people may be willing to accept as a group choice an option that is merely “close” to their ideal preferences.

Voting is a situation in which people may behave in this way. The usual starting model is a one-dimensional political spectrum, with conservative positions on the right and liberal positions on the left, as in Figure 1. We call each position on the spectrum a *platform* that a candidate or voter may choose to adopt. While a voter may represent her ideal platform by some point x on this line, she might be willing to vote for a candidate who is positioned at some point “close enough” to x , i.e., in an interval about x .




FIGURE 1. A one-dimensional political spectrum, with a single voter’s interval of approved platforms.

In this article, we ask the following: given such preferences on a political spectrum, when can we guarantee that some fraction (say, a majority) of the population will agree on some candidate? By “agree”, we mean in the sense of *approval voting*, in which voters declare which candidates they find acceptable.

Approval voting has not yet been adopted for political elections in the United States. However, many scientific and mathematical societies, such as the Mathematical Association of America and the American Mathematical Society, use approval voting for their elections. Additionally, countries other than the United States have used approval voting or an equivalent system; for details, see Brams and Fishburn [2] who discuss the advantages of approval voting.

Understanding which candidates can get voter approval can be helpful when there are a large number of candidates. An extreme example is the 2003 California gubernatorial recall election, which had 135 candidates in the mix [6]. We might imagine these candidates positioned at

*Corresponding author.

Research of various authors were partially supported by these NSF Grants: Berg by DMS-0301129, Norine by DMS-0200595, Su by DMS-0301129 and DMS-0701308, Thomas by DMS-0200595 and 0354742.

135 points on the line in Figure 1, which we think of as a subset of \mathbb{R} . If each California voter approves of candidates “close enough” to her ideal platform, we may ask under what conditions there is a candidate that wins the approval of a majority of the voters.

In this setting, we may assume that each voter’s set of approved platforms (her *approval set*) is a closed interval in \mathbb{R} , and that there is a set of candidates who take up positions at various points along this political spectrum. We shall call this spectrum with a collection of candidates and voters, together with voters’ approval sets, a *linear society* (a more precise definition will be given soon). We shall say that the linear society is *super-agreeable* if for every pair of voters there is some candidate that they would both approve, i.e., each pair of approval sets contains a candidate in their intersection. For linear societies this “local” condition guarantees a strong “global” property, namely, that there is a candidate that *every* voter approves! As we shall see in Theorem 5, this can be viewed as a consequence of Helly’s theorem about intersections of convex sets.

But perhaps this is too strong a conclusion. Is there a weaker local condition that would guarantee that only a majority (or some other fraction) of the voters would approve a particular candidate? For instance, we relax the condition above and call a linear society *agreeable* if among every three voters, some pair of voters approve the same candidate. Then it is not hard to show:

Theorem 1. *In an agreeable linear society, there is a candidate who has the approval of at least half the voters.*

More generally, call a linear society (k, m) -*agreeable* if it has at least m voters, and among every m voters, some subset of k voters approve the same candidate. Then our main theorem is a generalization of the previous result:

Theorem 2 (The Agreeable Linear Society Theorem). *Let $2 \leq k \leq m$. In a (k, m) -agreeable linear society, there is a candidate who has the approval of at least $(k-1)/(m-1)$ of the voters.*

We prove a slightly more general result in Theorem 8 and also briefly study societies whose approval sets are convex subsets of \mathbb{R}^d .

As an example, consider a city with fourteen restaurants along its main boulevard:

$A B C D E F G H I J K L M N$

and suppose every resident dines only at the five restaurants closest to his/her house (a set of consecutive restaurants, e.g., $DEFGH$). A consequence of Theorem 1 is that there must be a restaurant that is patronized by at least half the residents. Why? The pigeonhole principle guarantees that among every 3 residents, each choosing 5 of 14 restaurants, there must be a restaurant approved by at least 2 of them; hence this linear society is agreeable and Theorem 1 applies. For an example of Theorem 2, see Figure 3, which shows a $(2, 4)$ -agreeable linear society, and indeed there are candidates that receive at least $1/3$ of the votes (in this case $\lceil 7/3 \rceil = 3$).

We shall begin with some definitions, and explain connections to a classical convexity theorems, graph colorings, and maximal cliques in graphs. Then we prove Theorem 2, discuss extensions to higher-dimensional spectra, and conclude with some questions for further study.

2. DEFINITIONS

In this section, we fix terminology and explain the basic concepts upon which our results rely. Let us suppose that the set of all possible preferences is modeled by a set X , called the

spectrum. Each element of the spectrum is a *platform*. Assume that there is a finite set V of voters, and each voter v has an *approval set* A_v of platforms.

We define a *society* S to be a triple (X, V, \mathcal{A}) consisting of a spectrum X , a set of voters V , and a collection \mathcal{A} of approval sets for all the voters. Of particular interest to us will be the case of a *linear society*, in which X is a closed subset of \mathbb{R} and approval sets in \mathcal{A} are of the form $X \cap I$ where I is either empty or a closed bounded interval in \mathbb{R} . In general, however, X could be any set and approval sets could be any class of subsets of X . In Figure 2 we illustrate a linear society, where for ease of display we have separated the approval sets vertically so that they can be distinguished.




FIGURE 2. A linear society with infinite spectrum: each interval (shown here displaced above the spectrum) corresponds to the approval set of a voter. The shaded region indicates platforms with agreement number 4. This is a $(2, 3)$ -agreeable society.

Our motivation for considering intervals as approval sets arises from imagining that voters have an “ideal” platform along a linear scale (similar to Coombs’ J -scale [7]), and that voters are willing to approve “nearby” platforms, yielding approval sets that are connected intervals. Unlike the Coombs scaling theory, however, we are not concerned with preferential order of platforms; all platforms within a voter’s approval set have equivalent status as “approved” by that voter. We also note that while we model the spectrum as a linear scale, none of our results about linear societies depends on whether the scale is actually ordinal or cardinal.

We have seen that politics provides natural examples of linear societies. For a different example, X could represent a temperature scale, V a set of people that live in a house, and each A_v is a range of temperatures that person v finds comfortable. Then one may ask: at what temperature should the thermostat be set so as to satisfy the most number of people?




FIGURE 3. A linear society with a spectrum of two candidates (at platforms marked by carats): take the approval sets of the society of Figure 2 and intersect with these candidates. It is a $(2, 4)$ -agreeable linear society.

Two special cases of linear societies are worth mentioning. When $X = \mathbb{R}$ we should think of X as an infinite spectrum of platforms that potential candidates might adopt. However, in practice there are normally only finitely many candidates. We model that situation by letting

X be the set of platforms adopted by actual candidates. Thus one could think of X as either the set of all platforms, or the set of (platforms adopted by) candidates. See Figures 3 and 4.

Let $1 \leq k \leq m$ be integers. Call a society (k, m) -agreeable if it has at least m voters, and for any subset of m voters, there is at least one platform that at least k of them can agree upon, i.e., there is a point common to at least k of the voters' approval sets. Thus to be $(2, 3)$ -agreeable is the same as to be *agreeable*, and to be $(2, 2)$ -agreeable is the same as to be *super-agreeable*, as defined earlier.

One may check that the society of Figure 2 is $(2, 3)$ -agreeable. It is not $(3, 4)$ -agreeable, however, because among voters 1, 2, 4, 7 no three of them share a common platform. The same society, after restricting the spectrum to a set of candidates, is the linear society shown in Figure 3. It is not $(2, 3)$ -agreeable, because among voters 2, 4, 7 there is no pair that can agree on a candidate (in fact, voter 7 does not approve any candidate). However, one may verify that this linear society is $(2, 4)$ -agreeable.

For a society S , the *agreement number* of a platform, $a(p)$, is the number of voters in S who approve of platform p . The *agreement number* $a(S)$ of a society S is the maximum agreement number over all platforms in the spectrum, i.e.,

$$a(S) = \max_{p \in X} a(p).$$

The *agreement proportion* of S is simply the agreement number of S divided by the number of voters of S . This concept is useful when we are interested in percentages of the population rather than the number of voters. The society of Figure 2 has agreement number 4, which can be seen where the shaded rectangle covers the set that has maximum agreement number.

3. HELLY'S THEOREM AND SUPER-AGREEABLE SOCIETIES

Let us say that a society is \mathbb{R}^d -convex if the spectrum is \mathbb{R}^d and each approval set is a closed convex subset of \mathbb{R}^d . Note that an \mathbb{R}^1 -convex society is a linear society with spectrum \mathbb{R} . An \mathbb{R}^d -convex society can arise when considering a multi-dimensional spectrum, such as when evaluating political platforms over several axes (e.g., conservative vs. liberal, pacifist vs. militant, interventionist vs. isolationist). Or, the spectrum might be arrayed over more personal dimensions: the dating website *eHarmony* claims to use up to 29 of them [9]. In such situations, the convexity of approval sets might, for instance, follow from an independence-of-axes assumption and convexity of approval sets along each axis.

To find the agreement proportion of an \mathbb{R}^d -convex society, we turn to work concerning intersections of convex sets. The most well known result in this area is Helly's theorem. This theorem was proven by Helly in 1913, but the result was not published until 1921, by Radon [17].

Theorem 3 (Helly). *Given n convex sets in \mathbb{R}^d where $n > d$, if every $d + 1$ of them intersect at a common point, then they all intersect at a common point.*

Helly's theorem has a nice interpretation for \mathbb{R}^d -convex societies:

Corollary 4. *For every $d \geq 1$, a $(d + 1, d + 1)$ -agreeable \mathbb{R}^d -convex society must contain at least one platform that is approved by all voters.*

Notice that this theorem touts the existence of a common platform that lies in the intersection of all approval sets in a society. But with the introduction of candidates, there is no guarantee that there is a candidate in that common intersection. However, we can make such

a statement when $d = 1$ since the definition of a linear society includes the possibility that the spectrum is the collection of platforms adopted by a finite set of candidates.

Theorem 5 (The Super-Agreeable Linear Society Theorem). *A super-agreeable linear society must contain at least one platform that is approved by all voters.*

We provide a simple proof of this theorem, since the result will be needed later. When the spectrum is all of \mathbb{R} , this theorem is just Helly’s theorem for $d = 1$; a proof of Helly’s theorem for general d may be found in [15].

Proof. Let $X \subseteq \mathbb{R}$ denote the spectrum (recall that this could be a finite set of candidates or the entire real line). Since each voter v agrees on at least one platform with every other voter, we see that the approval sets A_v must be non-empty and contain a platform. Let $L_v = \min A_v$, $R_v = \max A_v$, and let $x = \max_v \{L_v\}$ and $y = \min_v \{R_v\}$. The first two minima/maxima exist because each A_v is compact; the last two exist because the number of voters is finite.




FIGURE 4. A super-agreeable linear society of 6 voters and 4 candidates, with agreement number 6.

We claim that $x \leq y$. Why? Since every pair of approval sets pairwise intersect in some platform, we see that $L_i \leq R_j$ for every pair of voters i, j . In particular, let i be the voter whose L_i is maximal and let j be the voter whose R_j is minimal. Hence $x \leq y$ and every approval set contains all platforms of X that are in the non-empty interval $[x, y]$, and in particular, the platform x . \square

The idea of this proof can be easily extended to furnish a proof of Theorem 1.

Proof. (Theorem 1) Using the same notations as in the prior proof, if $x \leq y$ then that proof shows that every approval set contains the platform x . Otherwise $x > y$ implies $L_i > R_j$ so that A_j and A_i do not contain a common platform.

We claim that for any other voter v , the approval set A_v contains either platform x or y (or both). After all, the society is agreeable so among the voters i, j, v some pair of A_i, A_j, A_v must contain a common platform; by the remarks above it must be that A_v intersects one of A_i or A_j . If A_v does not contain $x = L_i$ then since $L_v \leq L_i$ (by definition of x), we must have that $R_v < L_i$ and $A_v \cap A_i$ does not contain a platform. Then $A_v \cap A_j$ must contain a platform; hence $L_v \leq R_j$. Since $R_j \leq R_v$ (by definition of y), the platform $y = R_j$ must be in A_v .

Thus every approval set contains either x or y , and by the pigeonhole principle one of them must be contained in at least half the approval sets. \square

Proving the more general Theorem 2 will take a little more work.

4. THE AGREEMENT GRAPH OF LINEAR SOCIETIES IS PERFECT

To understand (k, m) -agreeability, it will be helpful to use a graph to represent the relation of intersecting approval sets. Recall that a *graph* G consists of a finite set $V(G)$ of *vertices* and a set $E(G)$ of 2-element subsets of $V(G)$, called *edges*. If $e = \{u, v\}$ is an edge, then we say that u, v are the *ends* of e , and that u and v are *adjacent* in G . We use uv as shorthand notation for the edge e .

Given a society S , we construct the *agreement graph* G of S by letting the vertices $V(G)$ be the voters of S and the edges $E(G)$ be all pairs of voters u, v whose approval sets intersect each other. Thus u and v are connected by an edge if there is a platform that both u, v would approve. Note that the agreement graph of society with agreement number equal to the number of voters is a complete graph (but the converse is false in higher dimensions, as we discuss later). Also note that a vertex v is isolated if A_v is empty or disjoint from other approval sets.




FIGURE 5. The agreement graph for the society in Figure 2. Note that voters 4, 5, 6, 7 form a maximal clique that corresponds to the maximal agreement number in Figure 2.

The *clique number* of G , written $\omega(G)$, is the greatest integer q such that G has a set of q pairwise adjacent vertices, called a *clique* of size q . By restricting our attention to members of a clique, and applying the Super-Agreeable Linear Society Theorem, we see that there is a platform that has the approval of every member of a clique, hence:

Fact 1. *For the agreement graph of a linear society, the clique number of the graph is the agreement number of the society.*

This fact does not necessarily hold if the society is not linear. For instance, it is easy to construct an \mathbb{R}^2 -convex society with three voters such that every two voters agree on a platform, but all three of them do not. It does, however hold in \mathbb{R}^d for *box societies*, to be discussed in Section 6.

Now, to get a handle on the clique number, we shall make a connection between the clique number and colorings of the agreement graph. The *chromatic number* of G , written $\chi(G)$, is the minimum number of colors necessary to color the vertices of G such that no two adjacent vertices have the same color. Thus two voters may have the same color as long as they do not agree on a platform. Note that in all cases, $\chi(G) \geq \omega(G)$.

A graph G is called an *interval graph* if every vertex x represents a closed interval $I_x \subseteq \mathbb{R}$ and $xy \in E(G)$ if and only if $I_x \cap I_y \neq \emptyset$. We have

Fact 2. *The agreement graph of a linear society is an interval graph.*

To see that Fact 2 holds let the linear society be (X, V, \mathcal{A}) , and let the voter approval sets be $A_v = X \cap I_v$, where I_v is a closed bounded interval or empty. We may assume that each I_v

is a minimal closed interval satisfying $A_v = X \cap I_v$; then the intervals $\{I_v : v \in V\}$ provide an interval representation of the agreement graph, as desired.

An *induced subgraph* of a graph G is a graph H such that $V(H) \subseteq V(G)$ and the edges of H are the edges of G that have both ends in $V(H)$. If every induced subgraph H of a graph G satisfies $\chi(H) = \omega(H)$, then G is called a *perfect graph*, e.g., see [18]. The following is a standard fact [20] about interval graphs:

Theorem 6. *Interval graphs are perfect.*

Proof. Let G be an interval graph, and for $v \in V(G)$, let I_v be the interval representing the vertex v . Since every induced subgraph of an interval graph is an interval graph, it suffices to show that $\chi(G) = \omega$, where $\omega = \omega(G)$. We proceed by induction on $|V(G)|$. The assertion holds for the null graph, and so we may assume that $|V(G)| \geq 1$, and that the statement holds for all smaller graphs. Let us select a vertex $v \in V(G)$ such that the right end of I_v is as small as possible. It follows that N , the set of neighbors of v in $V(G)$, are pairwise adjacent because their intervals must all contain the right end of I_v , and hence $|N| \leq \omega - 1$. See Figure 6. By induction, the graph $G \setminus v$ obtained from G by deleting v can be colored using ω colors, and since v has at most $\omega - 1$ neighbors, this coloring can be extended to a coloring of G , as desired. \square




FIGURE 6. If I_v, I_w intersect and the right end of I_v is smaller than the right end of I_w , then I_w must contain the right end of I_v .

The perfect graph property will allow us, in the next section, to make a crucial connection between the (k, m) -agreeability condition and the agreement number of the society. Given its importance in our setting, it is worth making a few comments about how perfect graphs appear in other contexts in mathematics, theoretical computer science, and operations research. The concept was introduced by Berge [1] in 1961, who was motivated by a question in communication theory, specifically, the determination of the Shannon capacity of a graph [19]. Chvátal later discovered that a certain class of linear programs always have an integral solution if and only if the corresponding matrix arises from a perfect graph in a specified way [5, 18, 3]. As pointed out in [18], algorithms to solve semi-definite programs grew out of the theory of perfect graphs. It has been proven recently [4] that a graph is perfect if and only if it has no induced subgraph isomorphic to a cycle of odd length at least five, or a complement of such a cycle.

5. (k, m) -AGREEABLE LINEAR SOCIETIES

We now use the connection between perfect graphs, the clique number, and the chromatic number to obtain a lower bound for the agreement number of a (k, m) -agreeable linear society (Theorem 8). We first need a lemma that says that in the corresponding agreement graph, the (k, m) -agreeable condition prevents any coloring of the graph from having too many vertices of the same color. Thus, there must be many colors and, since the graph is perfect, the clique number must be large as well.

Lemma 7. *Given integers $m \geq k \geq 2$, let positive integers q, ρ be defined by the division with remainder: $m - 1 = (k - 1)q + \rho$, where $0 \leq \rho \leq k - 2$. Let G be a graph on $n \geq m$ vertices with chromatic number χ such that every subset of $V(G)$ of size m includes a clique of size k . Then $n \leq \chi q + \rho$, or $\chi \geq (n - \rho)/q$.*

Proof. Let the graph be colored using the colors $1, 2, \dots, \chi$, and for $i = 1, 2, \dots, \chi$ let C_i be the set of vertices of G colored i . We may assume, by permuting the colors, that $|C_1| \geq |C_2| \geq \dots \geq |C_\chi|$. Since $C_1 \cup C_2 \cup \dots \cup C_{k-1}$ is colored using $k - 1$ colors, it includes no clique of size k , and hence, $|C_1 \cup C_2 \cup \dots \cup C_{k-1}| \leq m - 1$. It follows that $|C_{k-1}| \leq q$, for otherwise $|C_1 \cup C_2 \cup \dots \cup C_{k-1}| \geq (k - 1)(q + 1) \geq (k - 1)q + \rho + 1 = m$, a contradiction. Thus each $|C_i| \leq q$ for $i \geq k$ and

$$n = \sum_{i=1}^{k-1} |C_i| + \sum_{i=k}^{\chi} |C_i| \leq m - 1 + (\chi - k + 1)q = (k - 1)q + \rho + (\chi - k + 1)q = \chi q + \rho,$$

as desired. \square

Theorem 8. *Let $2 \leq k \leq m$. If G is the agreement graph of a (k, m) -agreeable linear society and q, ρ are defined by the division with remainder: $m - 1 = (k - 1)q + \rho$, $\rho \leq k - 2$, then the clique number satisfies:*

$$\omega(G) \geq \lceil (n - \rho)/q \rceil,$$

and this bound is best possible. It follows that this is also a lower bound on the agreement number, and hence every linear (k, m) -agreeable society has agreement proportion at least $(k - 1)/(m - 1)$.

Proof. By Fact 2 and Theorem 6 the graph G is perfect. Thus the chromatic number of G is equal to $\omega(G)$, and hence $\omega(G) \geq \lceil (n - \rho)/q \rceil$ by Lemma 7, as desired.

The second assertion follows from Fact 1 and noting that $(n - \rho)(m - 1) = n(k - 1)q + n\rho - \rho(m - 1) = n(k - 1)q + \rho(n - m + 1) \geq n(k - 1)q$, from which we see that $(n - \rho)/q \geq n(k - 1)/(m - 1)$.

Let us observe that the bound $\lceil (n - \rho)/q \rceil$ in Theorem 8 is best possible. Indeed, let I_1, I_2, \dots, I_q be disjoint intervals, for $i = q + 1, q + 2, \dots, n - \rho$ let $I_i = I_{i-q}$, and let $I_{n-\rho+1}, I_{n-\rho+2}, \dots, I_n$ be pairwise disjoint and disjoint from all the previous intervals, e.g., see Figure 7. Then the society with approval sets I_1, I_2, \dots, I_n is (k, m) -agreeable and its agreement graph has clique number $\lceil (n - \rho)/q \rceil$. \square




FIGURE 7. A linear $(4, 15)$ -society with $n = 21$ voters. Here $q = 4$, $\rho = 2$, so the clique number is at least $\lceil (n - \rho)/q \rceil = 5$.

The Agreeable Linear Society Theorem (Theorem 2) now follows as a corollary of Theorem 8.

6. \mathbb{R}^d -CONVEX AND d -BOX SOCIETIES

In this section we prove a higher dimensional analogue of Theorem 8 by giving a lower bound on the agreement proportion of a (k, m) -agreeable \mathbb{R}^d -convex society. We need a different method than our method for $d = 1$, because for $d \geq 2$, neither Fact 1 nor Fact 2 holds.

Also, we remark that, unlike linear societies, our results in this section about the agreement proportion for platforms will not necessarily hold when restricting the spectrum to a finite set of candidates in \mathbb{R}^d .

We shall use the following generalization of Helly's theorem, due to Kalai [13].

Theorem 9 (The Fractional Helly's Theorem). *Let $d \geq 1$ and $n \geq d + 1$ be integers, let $\alpha \in [0, 1]$ be a real number, and let $\beta = 1 - (1 - \alpha)^{1/(d+1)}$. Let F_1, F_2, \dots, F_n be convex sets in \mathbb{R}^d and assume that for at least $\alpha \binom{n}{d+1}$ of the $(d + 1)$ -element index sets $I \subseteq \{1, 2, \dots, n\}$ we have $\bigcap_{i \in I} F_i \neq \emptyset$. Then there exists a point in \mathbb{R}^d contained in at least βn of the sets F_1, F_2, \dots, F_n .*

The following is the promised analogue of Theorem 8.

Theorem 10. *Let $d \geq 1$, $k \geq 2$ and $m \geq k$ be integers. Then every (k, m) -agreeable \mathbb{R}^d -convex society has agreement proportion at least $1 - \left(1 - \binom{k}{d+1} / \binom{m}{d+1}\right)^{1/(d+1)}$.*

Proof. Let S be a (k, m) -agreeable \mathbb{R}^d -convex society, and let A_1, A_2, \dots, A_n be its voter approval sets. Let us call a set $I \subseteq \{1, 2, \dots, n\}$ good if $|I| = d + 1$ and $\bigcap_{i \in I} A_i \neq \emptyset$. By Theorem 9 it suffices to show that there are at least $\binom{k}{d+1} \binom{n}{d+1} / \binom{m}{d+1}$ good sets. We will do this by counting in two different ways the number N of all pairs (I, J) , where $I \subseteq J \subseteq \{1, 2, \dots, n\}$, I is good and $|J| = m$. Let g be the number of good sets. Since every good set is of size $d + 1$ and extends to an m -element subset of $\{1, 2, \dots, n\}$ in $\binom{n-d-1}{m-d-1}$ ways, we have $N = g \binom{n-d-1}{m-d-1}$. On the other hand, every m -element set $J \subseteq \{1, 2, \dots, n\}$ includes at least one k -element set K with $\bigcap_{i \in K} A_i \neq \emptyset$ (because S is (k, m) -agreeable), and K in turn includes $\binom{k}{d+1}$ good sets. Thus $N \geq \binom{k}{d+1} \binom{n}{m}$, and hence $g \geq \binom{k}{d+1} \binom{n}{d+1} / \binom{m}{d+1}$, as desired. \square

For $d = 1$, Theorem 10 gives a worse bound than Theorem 8, and hence for $d \geq 2$, the bound is most likely not best possible. However, a possible improvement must use a different method, because the bound in Theorem 9 is best possible.

A *box* in \mathbb{R}^d is the Cartesian product of d closed intervals, and we say that a society is a *d -box society* if each of its approval sets is a box in \mathbb{R}^d . By projection onto each axis, it follows from Theorem 5 that d -box societies satisfy the conclusion of Fact 1 (namely, that the clique number equals the agreement number), and hence their agreement graphs capture all the essential information about the society. Unfortunately, agreement graphs of d -box societies are, in general, not perfect. For instance, there is a 2-box society (Figure 8) whose agreement graph is the cycle on five vertices, hence its chromatic number is 3 but its clique number is 2.

For $k \leq m \leq 2k - 2$, the following theorem and corollary will resolve the agreement proportion problem for all (k, m) -agreeable societies satisfying the conclusion of Fact 1, and hence for all (k, m) -agreeable d -box societies where $d \geq 1$.

Theorem 11. *Let $m, k \geq 2$ be integers with $k \leq m \leq 2k - 2$, and let G be a graph on $n \geq m$ vertices such that every subset of $V(G)$ of size m includes a clique of size k . Then $\omega(G) \geq n - m + k$.*




FIGURE 8. A 2-box society whose agreement graph is a 5-cycle.

Before we embark on a proof let us make a few comments. First of all, the bound $n - m + k$ is best possible, as shown by the graph consisting of a clique of size $n - m + k$ and $m - k$ isolated vertices. Second, the conclusion $\omega(G) \geq n - m + k$ implies that every subset of $V(G)$ of size m includes a clique of size k , and so the two statements are equivalent under the hypothesis that $k \leq m \leq 2k - 2$. Finally, this hypothesis is necessary, because if $m \geq 2k - 1$, then for $n \geq 2(m - k + 1)$, the disjoint union of cliques of sizes $\lfloor n/2 \rfloor$ and $\lceil n/2 \rceil$ satisfies the hypothesis of Theorem 11, but not its conclusion.

A *vertex cover* of a graph G is a set $Z \subseteq V(G)$ such that every edge of G has at least one end in Z . We say a set $S \subseteq V(G)$ is *stable* if no edge of G has both ends in S . We deduce Theorem 11 from the following lemma.

Lemma 12. *Let G be a graph with minimum vertex cover of size z such that $G \setminus v$ has a vertex cover of size at most $z - 1$ for all $v \in V(G)$. Then $|V(G)| \leq 2z$.*

Proof. Let Z be a vertex cover of G of size z . For every $v \in V(G) - Z$ let Z_v be a vertex cover in $G \setminus v$ of size $z - 1$, and let $X_v = Z - Z_v$. Then X_v is a stable set. For $X \subseteq Z$ let $N(X)$ denote the set of neighbors of X outside Z . We have $v \in N(X_v)$ and $N(X_v) - \{v\} \subseteq Z_v - Z$, and so

$$|X_v| = |Z - Z_v| = |Z| - |Z \cap Z_v| = |Z_v| + 1 - |Z \cap Z_v| = |Z_v - Z| + 1 \geq |N(X_v)|.$$

On the other hand, if $X \subseteq Z$ is stable, then $|N(X)| \geq |X|$, for otherwise $(Z - X) \cup N(X)$ is a vertex cover in G of size at most $z - 1$, a contradiction. We have

$$(1) \quad |Z| \geq \left| \bigcup X_v \right| \geq \left| \bigcup N(X_v) \right| \geq |V(G)| - |Z|,$$

where both unions are over all $v \in V(G) - Z$, and hence $|V(G)| \leq 2z$, as required. To see that the second inequality holds let $u, v \in V(G) - Z$. Then

$$\begin{aligned} |X_u \cup X_v| &= |X_u| + |X_v| - |X_u \cap X_v| \geq |N(X_u)| + |N(X_v)| - |N(X_u \cap X_v)| \\ &\geq |N(X_u)| + |N(X_v)| - |N(X_u) \cap N(X_v)| = |N(X_u) \cup N(X_v)|, \end{aligned}$$

and, in general, the second inequality of (1) follows by induction on $|V(G) - Z|$. \square

Proof of Theorem 11. We proceed by induction on n . If $n = m$, then the conclusion certainly holds, and so we may assume that $n \geq m + 1$ and that the theorem holds for graphs on fewer than n vertices. We may assume that $m > k$, for otherwise the hypothesis implies

that G is the complete graph. We may also assume that G has two nonadjacent vertices, say x and y , for otherwise the conclusion holds. Then in G , every clique contains at most one of x, y , so in the graph $G \setminus \{x, y\}$ every set of vertices of size $m - 2$ includes a clique of size $k - 1$. Since $k - 1 \leq m - 2 \leq 2(k - 2) - 2$ we deduce by induction on n that $\omega(G) \geq \omega(G \setminus \{x, y\}) \geq n - 2 - (m - 2) + k - 1 = n - m + k - 1$. We may assume in the last statement that equality holds throughout, because otherwise G satisfies the conclusion of the theorem. Let \bar{G} denote the complement of G ; that is, the graph with vertex set $V(G)$ and edge set consisting of precisely those pairs of distinct vertices of G that are not adjacent in G . Let us notice that a set Q is a clique in G if and only if $V(G) - Q$ is a vertex cover in \bar{G} . Thus the size of a minimum vertex cover in \bar{G} is $m - k + 1$. Since $2(m - k + 1) \leq m \leq n$, by Lemma 12, the graph \bar{G} has an induced subgraph H on exactly m vertices with no vertex cover of size $m - k$ or smaller. By hypothesis, the graph \bar{H} has a clique Q of size k , but $V(H) - Q$ is a vertex cover in H of size $m - k$, a contradiction. \square

Theorem 13. *Let $d \geq 1$ and $m, k \geq 2$ be integers with $k \leq m \leq 2k - 2$, and let S be a (k, m) -agreeable d -box society with n voters. Then the agreement number of S is at least $n - m + k$, and this bound is best possible.*

Proof. The agreement graph G of S satisfies the hypothesis of Theorem 11, and hence it has a clique of size at least $n - m + k$ by that theorem. Since d -box societies satisfy the conclusion of Fact 1, the first assertion follows. The bound is best possible, because the graph consisting of a clique of size $n - m + k$ and $m - k$ isolated vertices is an interval graph. \square

7. DISCUSSION

As we have seen, set intersection theorems can provide a useful framework to model and understand the relationships between sets of preferences in voting, and this context leads to new mathematical questions. We suggest several directions which the reader may wish to explore.

Recent results in discrete geometry have social interpretations. The piercing number [11] of approval sets can be interpreted as the minimum number of platforms that are necessary such that everyone has some platform of which he or she approves. Set intersection theorems on other spaces (such as trees and cycles) are derived in [16] as an extension of both Helly's theorem and the KKM lemma [12]; as an application the authors show that in a super-agreeable society with a circular political spectrum, there must be a platform that has the approval of a strict majority of voters (in contrast with Theorem 5). Chris Hardin [10] has recently provided a generalization to (k, m) -agreeable societies with a circular political spectrum.

What results can be obtained for other spectra? The most natural problem seems to be to determine the agreement proportion for \mathbb{R}^d -convex and d -box (k, m) -agreeable societies. The smallest case where we do not know the answer is $d = 2$, $k = 2$, and $m = 3$. Rajneesh Hegde (private communication) found an example of a $(2, 3)$ -agreeable 2-box society with agreement proportion $3/8$. There may very well be a nice formula, because for every fixed integer d the agreement number of a d -box society can be computed in polynomial time. This is because the clique number of the corresponding agreement graph (also known as a graph of *boxicity* at most d) can be determined by an easy polynomial-time algorithm. On the other hand, for every $d \geq 2$ it is NP-hard to decide whether an input graph has boxicity at most d [14, 21]. (For $d = 1$ this is the same as testing whether a graph is an interval graph, and that can be done efficiently.)

Passing from results about platforms in societies to results about a finite set of candidates appears to be tricky in dimensions greater than 1. Are there techniques or additional hypotheses that would give useful results about the existence of candidates who have strong approval in societies with multi-dimensional spectra?

We may also question our assumptions. While convexity seems to be a rational assumption for approval sets in the linear case, in multiple dimensions additional considerations may become important. One might also explore the possibility of disconnected approval sets: what is the agreement proportion of a (k, m) -agreeable society in which every approval set has at most two components?

One might also consider varying levels of agreement. For instance, in a d -box society, two voters might not agree on every axis, so their approval sets do not intersect, but it might be the case that many of the projections of their approval sets do. In this case, one may wish to consider an agreement graph with weighted edges.

Finally, we might wonder about the agreement parameters k and m for various real-world issues. For instance, a society considering outlawing murder would probably be much more agreeable than that same society considering tax reform. Currently, we can empirically measure these parameters only by surveying large numbers of people about their preferences. It is interesting to speculate about methods for estimating suitable k and m from limited data.

This article grew out of the observation that Helly's theorem, a classical result in convex geometry, has an interesting voting interpretation. This led to the development of mathematical questions and theorems whose interpretations yield desirable conclusions in the voting context, e.g., Theorems 2, 8, 10, 13. It is nice to see that when classical theorems have interesting social interpretations, the social context can also motivate the study of new mathematical questions.

8. ACKNOWLEDGEMENTS

The authors wish to thank a referee for many valuable suggestions.

REFERENCES

- [1] C. Berge. Färbung von Graphen deren sämtliche bzw. deren ungerade Kreise starr sind (Zusammenfassung), *Wissenschaftliche Zeitschrift*, Martin Luther Universität Halle-Wittenberg, Mathematisch-Naturwissenschaftliche Reihe (1961), 114–115.
- [2] S. J. Brams and P. C. Fishburn. Going from theory to practice: the mixed success of approval voting, *Social Choice and Welfare* **25** (2005), 457–474.
- [3] M. Chudnovsky, N. Robertson, P. D. Seymour and R. Thomas. Progress on perfect graphs, *Math. Program. Ser. B* **97** (2003), 405–422.
- [4] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas. The strong perfect graph theorem, *Ann. Math.* **164** (2006), 51–229.
- [5] V. Chvátal. On certain polytopes associated with graphs, *J. Combin. Theory Ser. B* **13** (1975), 138–154.
- [6] CNN. California ballot official, ready to print. CNN.com, August 14, 2003. <<http://www.cnn.com/2003/ALLPOLITICS/08/13/calif.recall/index.html>>, October 2008.
- [7] C.H. Coombs. *A Theory of Data*. New York: John Wiley and Sons, 1964.
- [8] B. Disraeli. From *The Columbia World of Quotations*. New York: Columbia University Press, 1996. Also online at <<http://www.bartleby.com/66/88/16788.html>>, October 2008.
- [9] eHarmony.com. What are the 29 Dimensions? <<http://www.eharmony.com/why/dimensions>>, October 2008.
- [10] C. Hardin. Agreement in circular societies, preprint, 2007.
- [11] M. Katchalski and D. Nashtir. A Helly type conjecture. *Discrete Comput. Geom.*, 21(1):37–43, 1999.
- [12] B. Knaster, C. Kuratowski, and S. Mazurkiewicz. Ein Beweis des Fixpunktsatzes für n -dimensionale Simplexe. *Fund. Math.*, 14:132–137, 1929.
- [13] G. Kalai. Intersection patterns of convex sets, *Israel J. Math.*, 48:161–174, 1984.

- [14] J. Kratochvíl. A special planar satisfiability problem and some consequences of its NP-completeness, *Discrete Appl. Math.* **52** (1994), 233-252.
- [15] J. Matousek. *Lectures on Discrete Geometry*. New York: Springer-Verlag, 2002.
- [16] A. Niedermaier, D. Rizzolo, and F. E. Su. A tree segmentation theorem, preprint 2006.
- [17] J. Radon. Mengen konvexer Körper, die einen gemeinsamen Punkt enthalten, *Math. Ann.* **83** (1921), 113–115.
- [18] J. L. Ramírez Alfonsín and B. A. Reed. *Perfect graphs*. Chichester: J. Wiley and Sons, 2001.
- [19] C. E. Shannon. The zero error capacity of a noisy channel, *I.R.E. Trans. Inform. Theory IT-2* (1956), 8–19.
- [20] D.B. West. *Introduction to Graph Theory*. Upper Saddle River, NJ: Prentice-Hall, 2001.
- [21] M. Yannakakis. The complexity of the partial order dimension problem, *SIAM J. Algebraic Discrete Methods* **3** (1982), 351–358.

This material is based upon work supported by the National Science Foundation under Grants No. 0200595 and 0354742 and 0301129 and 0701308. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA, LINCOLN NE 68588, debbie.berg@gmail.com

DEPARTMENT OF MATHEMATICS, FINE HALL, WASHINGTON ROAD, PRINCETON NJ, 08544, snorin@math.princeton.edu

DEPARTMENT OF MATHEMATICS, HARVEY MUDD COLLEGE, CLAREMONT CA 91711, su@math.hmc.edu

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332, thomas@math.gatech.edu

MATHEMATISCHES SEMINAR DER UNIVERSITÄT HAMBURG, BUNDESSTR. 55, D-20146 HAMBURG GERMANY, wollan@math.uni-hamburg.de